

# Fluarix Tetra

## Quadrivalent influenza vaccine (split virion, inactivated)

### 1. NAME OF THE MEDICINAL PRODUCT

Fluarix Tetra suspension for injection in pre-filled syringe  
Influenza vaccine (split virion, inactivated)

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Fluarix Tetra is an inactivated influenza vaccine (split virion), containing antigens (propagated in embryonated eggs) equivalent to the following types and subtypes:

A/Michigan/45/2015 (H1N1)pdm09 - like strain (A/Singapore/GP1908/2015, IVR-180);

A/Singapore/INFIMH-16-0019/2016 (H3N2) – like strain (A/Singapore/INFIMH-16-0019/2016, NIB-104);

B/Brisbane/60/2008 - like strain (B/Brisbane/60/2008, wild type);

B/Phuket/3073/2013 - like strain (B/Phuket/3073/2013, wild type).

Each 0.5 ml vaccine dose contains 15 µg haemagglutinin of each of the recommended strains.

This vaccine complies with the World Health Organisation (WHO) recommendation (Southern Hemisphere) for the **2018** season.

For the full list of excipients see *section 6.1*.

Fluarix Tetra may contain traces of Formaldehyde, Sodium Deoxycholate, Ovalbumin, Gentamicin Sulphate and Hydrocortisone. The maximum amount of Ovalbumin that may be present is not more than 0.05 micrograms per dose.

### 3. PHARMACEUTICAL FORM

Suspension for injection in pre-filled syringe.  
The suspension is colourless and slightly opalescent.

### 4. CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

Fluarix Tetra is indicated for active immunisation of adults and children from 3 years of age for the prevention of influenza disease caused by the two influenza A virus subtypes and the two influenza B virus types contained in the vaccine.

The use of Fluarix Tetra should be based on official recommendations.

## **4.2 Posology and method of administration**

### Posology

Adults: 0.5 ml

### *Paediatric population*

Children from 36 months onwards: 0.5 ml.

For children aged < 9 years, who have not previously been vaccinated, a second dose should be given after an interval of at least 4 weeks.

Vaccination should be carried out by intramuscular injection preferably into the deltoid muscle or anterolateral thigh (depending on the muscle mass).

Children less than 3 years: the safety and efficacy of Fluarix Tetra in children less than 3 years have not been established.

### Method of administration

Immunisation should be carried out by intramuscular injection.

Precautions to be taken before handling or administering the medicinal product.

For instructions for preparation of the medicinal product before administration, see *section 6.6*.

## **4.3 Contraindications**

Hypersensitivity to the active substances or to any of the excipients listed in section 6.1 or to any component that may be present as traces such as eggs (ovalbumin, chicken proteins), formaldehyde, gentamicin sulphate and sodium deoxycholate.

Immunisation should be postponed in patients with febrile illness or acute infection.

## **4.4 Special warnings and precautions for use**

It is good clinical practice to precede vaccination by a review of the medical history (especially with regard to previous vaccination and possible occurrence of undesirable events) and a clinical examination.

As with all injectable vaccines, appropriate medical treatment and supervision should always be readily available in case of an anaphylactic event following the administration of the vaccine.

Antibody response in patients with endogenous or iatrogenic immunosuppression may be insufficient.

Fluarix Tetra is not effective against all possible strains of influenza virus. Fluarix Tetra is intended to provide protection against those strains of virus from which the vaccine is prepared and to closely related strains.

As with any vaccine, a protective immune response may not be elicited in all vaccinees.

Fluarix Tetra should under no circumstances be administered intravascularly.

As with other vaccines administered intramuscularly, Fluarix Tetra should be given with caution to individuals with thrombocytopenia or any coagulation disorder since bleeding may occur following an intramuscular administration to these subjects.

Syncope (fainting) can occur following, or even before, any vaccination especially in adolescents as a psychogenic response to the needle injection. This can be accompanied by several neurological signs such as transient visual disturbance, paraesthesia and tonic-clonic limb movements during recovery. It is important that procedures are in place to avoid injury from faints.

Interference with serological testing.

See *section 4.5*.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

No interaction studies have been performed. If Fluarix Tetra is to be given at the same time as another injectable vaccine, the vaccines should always be administered at different injection sites.

Following influenza vaccination, false-positive results in serology tests using the ELISA method to detect antibodies against HIV1, Hepatitis C and especially HTLV1 have been observed. The Western Blot technique disproves the false-positive ELISA test results. The transient false-positive reactions could be due to the IgM response by the vaccine.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

Inactivated influenza vaccines can be used in all stages of pregnancy. Larger datasets on safety are available for the second and third trimester, compared with the first trimester; however, data from worldwide use of inactivated influenza vaccines do not indicate any adverse foetal and maternal outcomes attributable to the vaccine (see *Pre-clinical Safety Data*).

##### Breast-feeding

Fluarix Tetra may be used during breast-feeding.

##### Fertility

No fertility data are available.

#### **4.7 Effects on ability to drive and use machines**

Fluarix Tetra has no or negligible influence on the ability to drive and use machines.

#### **4.8 Undesirable effects**

In two clinical studies, healthy adults 18 years of age and older and healthy children 3 to 17 years of age were administered Fluarix Tetra (more than 3,000 adults and 900 children) or GlaxoSmithKline trivalent influenza vaccine, Fluarix (more than 1,000 adults and 900 children). Similar rates of solicited adverse events were observed in recipients of Fluarix Tetra and Fluarix.

Adverse reactions reported for Fluarix Tetra are listed per dose according to the following frequency categories:

Very common ( $\geq 1/10$ );  
 Common ( $\geq 1/100$  to  $< 1/10$ );  
 Uncommon ( $\geq 1/1,000$  to  $< 1/100$ );  
 Rare ( $\geq 1/10,000$  to  $< 1/1,000$ );  
 Very rare ( $< 1/10,000$ )

Frequency	Adverse reactions
<b>Clinical trials</b>	
Very common	irritability <sup>1</sup> , myalgia, injection site pain, fatigue
Common	appetite loss <sup>1</sup> , drowsiness <sup>1</sup> , headache, gastrointestinal symptoms (including nausea, vomiting, diarrhoea and/or abdominal pain), arthralgia, injection site redness <sup>4</sup> , injection site swelling <sup>4</sup> , shivering, fever
Uncommon	dizziness <sup>2</sup> , rash <sup>3</sup> , injection site hematoma <sup>2</sup> , injection site pruritus
<sup>1</sup> reported as a solicited symptom in subjects less than 6 years of age <sup>2</sup> reported in adult subjects <sup>3</sup> reported in subjects 3 years to 17 years of age <sup>4</sup> very common in subjects 3 years to 17 years of age	
<b>Adverse reactions additionally reported in previous Fluarix trials:</b>	
Common	Sweating, injection site induration
<b>Post-marketing experience</b>	
As all three of the influenza strains contained in Fluarix are included in Fluarix Tetra, the following adverse events that have been observed for Fluarix during post-marketing surveillance may occur in patients receiving Fluarix Tetra post-approval.	
Rare	transient lymphadenopathy, allergic reactions (including anaphylactic reactions), neuritis, acute disseminated encephalomyelitis, Guillain-Barré syndrome*, urticaria, pruritus, erythema, angioedema, influenza-like illness, malaise  *Spontaneous reports of Guillain-Barré syndrome have been received following vaccination with Fluarix; however, a causal association between vaccination and Guillain-Barré syndrome has not been established.

#### 4.9 Overdose

Insufficient data are available.

### 5. PHARMACOLOGICAL PROPERTIES

#### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Influenza vaccine, ATC Code: J07BB02

#### Mechanism of action

Fluarix Tetra provides active immunisation against four influenza virus strains (two A subtypes and two B types) contained in the vaccine and induces humoral antibodies against the haemagglutinins. These antibodies neutralise influenza viruses.

Specific levels of hemagglutination-inhibition (HI) antibody titer post-vaccination with inactivated influenza virus vaccines have not been correlated with protection from influenza illness but the HI antibody titers have been used as a measure of vaccine activity. In some human challenge studies, HI antibody titres of  $\geq 1:40$  have been associated with protection from influenza illness in up to 50% of subjects.

#### Pharmacodynamic effects

#### **Immunogenicity of Fluarix Tetra versus Fluarix**

Clinical studies performed in adults (D-QIV-001 and D-QIV-008) and in children 3 years to 17 years of age (D-QIV-003) assessed the non-inferiority of Fluarix Tetra versus Fluarix for HI Geometric mean antibody titer (GMT) at Day 21 (for adults) and at Day 28 (for children) and HI seroconversion rate (4-fold rise in reciprocal titer or change from undetectable [ $< 10$ ] to a reciprocal titer of  $\geq 40$ ).

In all studies, the immune response elicited by Fluarix Tetra against the three strains in common was non-inferior to Fluarix. Fluarix Tetra elicited a superior immune response against the additional B strain included in Fluarix Tetra compared to Fluarix.

#### Adults 18 years of age and older

In clinical study D-QIV-008, approximately 1,800 adults 18 years of age and older received a single dose of Fluarix Tetra and approximately 600 subjects received a single dose of Fluarix (Table 1).

**Table 1: Post-vaccination GMT and seroconversion rates:**

<b>Adults 18 years of age and older</b>	<b>Fluarix Tetra N=1809</b>	<b>Fluarix<sup>1</sup> N=608</b>
	<b>GMT (95% confidence interval)</b>	
<b>A/H1N1</b>	201.1 (188.1;215.1)	218.4 (194.2;245.6)
<b>A/H3N2</b>	314.7 (296.8;333.6)	298.2 (268.4;331.3)
<b>B (Victoria)<sup>2</sup></b>	404.6 (386.6;423.4)	393.8 (362.7;427.6)
<b>B (Yamagata)<sup>3</sup></b>	601.8 (573.3;631.6)	386.6 (351.5;425.3)
	<b>Seroconversion rate (95% confidence interval)</b>	
<b>A/H1N1</b>	77.5% (75.5;79.4)	77.2% (73.6;80.5)
<b>A/H3N2</b>	71.5% (69.3;73.5)	65.8% (61.9;69.6)
<b>B (Victoria)</b>	58.1% (55.8;60.4)	55.4% (51.3;59.4)
<b>B (Yamagata)</b>	61.7% (59.5;64.0)	45.6% (41.6;49.7)

<sup>1</sup>containing A/H1N1, A/H3N2 and B (Victoria lineage)

<sup>2</sup>recommended strain by WHO during the season 2010-2011

<sup>3</sup>additional B strain contained in Fluarix Tetra recommended in season 2008-2009

Post-vaccination seroprotection rates (Day 21 reciprocal titer of  $\geq 40$ ) for Fluarix Tetra were 91.3% against A/H1N1, 96.8% against A/H3N2, 98.8% against B (Victoria) and 91.8% against B (Yamagata).

In clinical study D-QIV-001 (vaccine composition of 2007-2008 season), post-vaccination seroprotection rates for Fluarix Tetra were 92.3% against A/H1N1, 97.1% against A/H3N2, 97.1% against B (Victoria) and 98.1% against B (Yamagata).

#### Children 3-17 years of age

In clinical study (D-QIV-003), approximately 900 children from 3-17 years of age received one or two doses of Fluarix Tetra or Fluarix, respectively (Table 2).

**Table 2: Post-vaccination GMT and seroconversion rates:**

Children 3 years to 17 years of age	Fluarix Tetra N=791	Fluarix <sup>1</sup> N=818
	<b>GMT (95% confidence interval)</b>	
<b>A/H1N1</b>	386.2 (357.3;417.4)	433.2 (401.0;468.0)
<b>A/H3N2</b>	228.8 (215.0;243.4)	227.3 (213.3;242.3)
<b>B (Victoria)<sup>2</sup></b>	244.2 (227.5;262.1)	245.6 (229.2;263.2)
<b>B (Yamagata)<sup>3</sup></b>	569.6 (533.6;608.1)	224.7 (207.9;242.9)
	<b>Seroconversion rate (95% confidence interval)</b>	
<b>A/H1N1</b>	91.4% (89.2;93.3)	89.9% (87.6;91.8)
<b>A/H3N2</b>	72.3% (69.0;75.4)	70.7% (67.4;73.8)
<b>B (Victoria)</b>	70.0% (66.7;73.2)	68.5% (65.2;71.6)
<b>B (Yamagata)</b>	72.5% (69.3;75.6)	37.0% (33.7;40.5)

<sup>1</sup>containing A/H1N1, A/H3N2 and B (Victoria lineage)

<sup>2</sup>recommended strain by WHO during the season 2010-2011

<sup>3</sup>additional B strain contained in Fluarix Tetra recommended in season 2008-2009

Post-vaccination seroprotection rates for Fluarix Tetra were 96.6% against A/H1N1, 98.0% against A/H3N2, 97.3% against B (Victoria) and 99.2% against B (Yamagata).

## **5.2 Preclinical safety data**

Non-clinical data reveal no special hazards for humans based on conventional studies of acute toxicity, local tolerance, repeated dose toxicity and reproductive/developmental toxicity.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Sodium chloride, disodium phosphate dodecahydrate, potassium dihydrogen phosphate, potassium chloride, magnesium chloride hexahydrate,  $\alpha$ -tocopheryl hydrogen succinate, polysorbate 80, octoxinol 10 and water for injections.

Hydrocortisone, gentamicin sulfate, ovalbumin, formaldehyde and sodium deoxycholate are present as residuals from the manufacturing process.

### **6.2 Incompatibilities**

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

The expiry date is indicated on the label and packaging.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C to 8°C).

Do not freeze.

Store in the original package in order to protect from light.

### **6.5 Nature and contents of the container**

0.5 ml suspension in prefilled syringe (Type I glass) with a plunger stopper (grey butyl rubber) with fixed or separate or without needles in the following pack sizes:

- with fixed needle: pack sizes of 1 or 10
- with 1 separate needle: pack sizes of 1 or 10
- with 2 separate needles: pack size of 1
- without needle: pack sizes of 1 or 10

The needle type(s) that may be supplied in the pack are: 23G 1", 25G 5/8" and 25G 1".

Not all presentations are available in every country.

### **6.6 Special precautions for disposal and other handling**

The vaccine should be allowed to reach room temperature before use.


The vaccine presents as a colourless to slightly opalescent suspension.

The syringe should be shaken and inspected visually for any foreign particulate matter and/or variation of physical aspect prior to administration. In the event of either being observed, discard the vaccine.



#### Instructions for administration of the vaccine presented in pre-filled syringe without a fixed needle

To attach the needle to the syringe, refer to the below drawing. However, the syringe provided with Fluarix Tetra might be slightly different (without screw thread) than the syringe described in the drawing. In that case, the needle should be attached without screwing.

## Needle



## Syringe



1. Holding the syringe **barrel** in one hand (avoid holding the syringe plunger), unscrew the syringe cap by twisting it anticlockwise.
2. To attach the needle to the syringe, twist the needle clockwise into the syringe until you feel it lock. (see picture)
3. Remove the needle protector, which on occasion can be a little stiff.
4. Administer the vaccine.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Trade marks are owned by or licensed to the GSK group of companies.

Version number: GDS01/IPI08SI (SH)

Date of issue: November 2017

© 2017 GSK group of companies or its licensor

Manufacturer:

GlaxoSmithKline Biologicals, Branch of SmithKline Beecham Pharma GmbH & Co. KG,  
Dresden, Germany

<GSK logo>